



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ished, some of this imprisoned air escapes, and the greater the fall of the barometer the greater the force with which the air is expelled. My friend, Mr. Smith, utilized this air current to blow a whistle, which could be heard all over the town, warning the inhabitants of a possible storm. With a rising barometer, caused by an increase in the pressure of the air, air will be forced back into the subterranean reservoir. Mr. Smith tells me that when the air is going into the well, the water recedes a certain amount, and that when the air is blowing out, it can be heard bubbling through the water.

ADDITIONAL NOTE ON THE BRENHAM METEORITE.

BY ROBERT HAY.

About the end of 1891, the finds of the meteoric fall in Kiowa county were extended nearly a mile east of the former ones, and most of them are of a new type. Several groups were found, each in an area of several square yards, and having several hundred individuals. The aggregate number was about 3,000. Some of them




FIG. 14. POLISHED SECTION OF BRENHAM METEORITE; *a*, cavities containing olivene; *b*, Widmanstätten figures on polished surface.

seemed to be the decomposed parts of a larger mass, but the bulk of them were evidently separate meteorites. A few were about a pound in weight; others were from six or seven ounces down to the size of a pea. All were more or less oxidized; some had lost all their metallic structure, but some, even of the very smallest, had the true pallasite structure. A specimen (exhibited to the Academy) had been pronounced by Professor Foote, of Philadelphia, to be almost identical with the original meteorite of Pallas, which gives the name to this variety. There was one mass of nearly 80 pounds. There seem to have been no more finds, though the search was active.

NOTE ON THE OCCURRENCE OF GRANITE IN A DEEP BORING IN EASTERN KANSAS.

BY ROBERT HAY, F. G. S. A.

Four borings (one reaching 1,000 feet in depth) at Fort Scott have passed through the coal measures and subcarboniferous rocks at that place. The deep boring at Pittsburg (1,200 feet) is said by Mr. St. John to reach silurian rocks. The boring at Leavenworth (1,800 feet) is also said to have its bottom in siluria. Neither the well seven miles east of Wichita (1,943 feet) nor the boring at Anthony (2,300 feet)